Document Title: Q-Pulse Index:

Revision Number: [redacted] **Author:** [redacted]

Hepatitis C Viral Load 49

Hepatitis D (Delta agent) 49

[illegible] ..... [illegible]

50 Herpes Simplex Virus 1 and 2 PCR (Qualitative) ..... [illegible]

Nose Swab Culture.....63
Parasitology Investigation.....63
Parasitology investigation for *Enterobius vermicularis*.....63

Document Title: Q-Pulse Index:

Revision Number: [redacted] **Author:** [redacted]

We advise all healthcare workers NOT to ask for results pertaining to themselves, but to obtain test results from the requesting physician, their doctor or from Occupational Health as appropriate.

We request that users do not phone the lab to confirm whether samples have been sent or not, as this takes up much valuable time and prevents lab staff from completing their work in a timely fashion. We recommend that patient notes are annotated to confirm samples requested and taken.

4.6. Telephoned results

Results of urgent requests and results which may aid the immediate patient management will be telephoned. This includes all positive blood cultures, positive CSFs, and other clinically significant results.

All other results will only be telephoned on request.

4.7. Turnaround times

The laboratory continually monitors its turnaround times to ensure that it complies with its responsibilities within the patient pathway. The laboratory measures its turnaround times as the time from receipt until the point at which the result is authorised.

The expected turnaround times for each test are indicated on the individual test sheets. For detailed turnaround times for each test and actual performance, please contact the laboratory.

Interrogation of the electronic systems allows for full audit of the reception, testing and reporting process, including time of report viewing and report printing.

5. Sample Collection

5.1. Viral and bacterial serology tests

As a general guide, a 4mL yellow top vacutainer tube is adequate for up to three viral serology screening tests plus provide sufficient sample to be used for referral to the reference laboratory if the screening test is positive.

For four or more tests, two 4mL samples are advised. For unusual
performed at SDH, an additional sample is advised to speed up handling and packaging.

Requests received on Laboratory Medicine (blue) request forms will NOT be accepted. Please use only the appropriate request on T-quest the or Microbiology (black) request forms for viral & bacterial serology tests.

5.2. Optimum time of and conditions for collection

Samples for bacterial culture, wherever possible, should be collected prior to commencement of antibiotic treatment.

Actual pus or tissue samples are always preferable to a swab.

Document Title: Q-Pulse Index:

clearly labelled. Once collected, place the specimen into a plastic specimen bag and seal the bag

	Bacteriology swabs in Amies transport medium with charcoal	Materials Management Team
	Bacteriology swabs in Amies transport medium	Materials Management Team
	Pernasal swab for whooping cough	Materials Management Team
	Viral swabs in viral transport medium	Materials Management Team
	Faeces container	Materials Management Team

Sterile univ 15 6.46 36.12 reW*nBT/1

- Tuberculosis (samples from sites where tuberculosis infection is likely)
- Anthrax
- Brucellosis
- Transmissible Spongiform Encephalopathy (including CJD)
- Viral haemorrhagic fever
- Avian Flu
- MERS

Please refer to the T r u s Pd l i c y fs o r the Transport of Pathology Specimens.

To minimise the risks, ensure that such samples are packaged as follows:

- A t t ach a " Da n ge r o f In fec t i o n " l abe l t o t he s amp l e co n t ain e r a n d r eq u
  qualifying samples (including biochemistry and haematology requests)7.002 (ensu)4.004 (re)-2.002 (that)9.0

Document Title: Q-Pulse Index:

Revision Number: [redacted] **Author:** [redacted]

Type of Clinical Material	Retention time
[redacted]	[redacted]
[redacted]	[redacted]
[redacted]	[redacted]

Unless stated otherwise in the test table, please send at least 200µL of

	Ideally samples should be collected before antibiotic treatment. Delays in transportation may affect the recovery of pathogens.
	No

Atypical Pneumonia CFTs

	Atypical Pneumonia CFTs This test includes: • Influenza A, • Influenza B, • Respiratory syncytial virus (RSV), • *Chlamydia* species, • *Mycoplasma pneumoniae*, • Q Fever Phase II, • Adenovirus
	Complement fixation test (CFT)
	Clotted blood
	Yellow top (minimum volume 3.5 ml)
	The timing of the onset of patient symptoms and the blood sample(s) taken is critical. Samples taken less than 10 days after onset of symptoms is considered an ACUTE sample and will be stored pending arrival of a CONVALESCENT sample (taken 10 t

	X 2 EDTA blood samples
	Purple top (minimum volume 500µl)
	No special requirements Specimens should be sent to the laboratory without delay during normal working hours. Outside of normal working hours samples should be refrigerated.
	10-14 days
	For non-renal patients, or not specified to go to QA Hospital Portsmouth.
	No
	This test is performed at UKHSA Bristol Reference Laboratory.

Blood culture

Blood cultures are used to detect the cause of an infection leading to bacteraemia or fungaemia. The results are important because they help guide appropriate treatment.
Results of all significant positive blood cultures will be telephoned to a clinician as soon as they become positive. As the isolation time depends upon the organism and the initial inoculum, this may vary from a few hours, up to five days after receipt.

	Blood Cultures
	Gram stain, if positive & Culture
	Adult –5-10 ml of blood per bottle. Paediatric –3-4 ml of blood.
	Adult blood culture set –Aerobic (blue) and Anaerobic (purple) bottles. Paediatric blood culture bottle –yellow top.

	False negatives may occur if inadequate blood culture volumes are submitted. Do not remove or cover up barcode labels as these are required in the laboratory.
	Bottles should be left at room temperature in blood-issue room.

Bordetella pertussis Culture

Whooping cough is a highly contagious disease that is caused by the fastidious Gram-negative coccobacillus *Bordetella pertussis*. In some cases, this syndrome may also be caused by *Mycoplasma pneumoniae*, and by viruses such as adenoviruses and enteroviruses.

	Bordetella pertussis culture
	Isolation and characterisation of *Bordetella* species.
	Pernasal swab
	Pernasal swab (blue top)

	Pertussis serology is usually more useful in adults presenting with a prolonged cough. Please state onset date. Vaccination history may help interpretation of results.
	No
	This test is performed at UKHSA Bristol Reference Laboratory.

Bordetella pertussis PCR

Bordetella pertussis

ijoux containlastic

	Cell count, Gram stain and Culture
	1 –2ml CSF.

Two sterile plastic bijoux containers.

	The laboratory now screens for *Neisseria gonorrhoeae* for both hospital and community patients. If you do NOT wish to have *N. gonorrhoeae* tested on individual patients, please make this clear on the request form (in the clinical details box).

Chlamydia trachomatis Infection (genital swabs)

	Detection of *Chlamydia trachomatis* DNA in genital swabs
	PCR
	• Endo-cervical swab, • HVS, • Vulvo-vaginal swabs.
	Cobas PCR female swab kit (Yellow Top)
	Specimens should be collected and handled following the recommended guidelines on the collection kits. Illustrated collection procedure is provided on the back of the collection kits. Specimens should be sent to the laboratory without delay during normal working hours.
	7 days
	Clinical details are essential for processing.
	No
	Chlamydia swabs have TWO separate expiry dates: one for the swab and one for the transport media contained in the pack. Note that the expiry date of the swab may differ by some months to that of the transport media. It is usually the media which has the shortest expiry date. On the Cobas PCR chlamydia swab, the expiry date can be found at the bottom of the blister pack, below the Lot number on the pack. The date is printed in the reverse order to that we normally use in the UK, i.e., YEAR/ MONTH, so March 2023 would appear as 2023/03. Please return any out-of-date swabs to the Microbiology Laboratory and request replacements as required. Please ensure that the lid of the Cobas PCR tube is securely tightened to prevent sample from leaking. The laboratory now screens for *Neisseria gonorrhoeae* for both hospital and community patients. IF you do NOT wish to have *N. gonorrhoeae* tested on individual patients, please make this clear on the request form (in the clinical details box).

Chlamydia trachomatis infection (Non-genital sites: Eye, Throat, Rectum)

	Detection of *Chlamydia trachomatis* DNA in non-genital swabs (Eye, Throat, Rectum, etc.)
	PCR
	Swab from appropriate site
	Cobas PCR female swab sample pack (Yellow Top)

Clostridium difficile Toxin detection

C. difficile is a Gram positive, spore forming, strictly anaerobic rod, so named because of the difficulty in original culture and characterisation. Toxigenic strains produce large protein toxins A and B, both being major virulence factors. Most disease associated with *C. difficile* is intestinal, though *C. difficile* may be isolated from blood or tissues. Changes in the gut flora associated with broad spectrum antibiotics and chemotherapeutic agents can result in colonisation by *C. difficile*.

	Clostridium difficile Toxin
	Toxin Detection Faeces

	Specimens should be sent to the laboratory without delay during normal working hours. Outside of normal working hours samples should be refrigerated.
	10-14 days
	Clinical details are essential for processing.
	No
	This test is processed at UKHSA Bristol Reference Laboratory.

Viral swabs in viral transport medium

	No This tes

	Purple top (minimum volume 4 ml)
	No special requirements. Specimens should be sent to the laboratory without delay during normal working hours. Outside of normal working hours samples should be refrigerated. 10-14 days

	Specimens should be sent to the laboratory without delay during normal working hours. Outside of normal working hours samples should be refrigerated.
	10-14 days
	No
	This test is processed at UKHSA Bristol Reference Laboratory.

Full Respiratory Panel (PCR)

	Detection of respiratory viruses
	PCR
	Viral swab
	Green topped swab Red topped swab
	No special requirements. Specimens should be sent to the laboratory without delay during normal working hours. Outside of normal working hours samples should be refrigerated.

H. pylori stool antigen testing is more specific and allows post treatment testing or re-testing if symptoms re-occur despite therapy.
The alternative test (if faeces is unacceptable to the patient) is the Urea Breathe Test. This should be arranged through the Gastroenterology Department or via Community prescription at the local pharmacy.
Public Health England produces a useful guide on who and when to test for *H. pylori*

Document Title:_ O-Pulse Index:

Revision Number:

	This test is performed at UKHSA Bristol Reference Laboratory.

Hepatitis C Viral Load

	Hepatitis C Viral Load
	PCR
	2 x EDTA blood
	Purple top (minimum volume 4 ml)
	No special requirements. Specimens should be sent to the laboratory without delay during normal working hours. Outside of normal working hours samples should be refrigerated.
	10-14 days
	Requests must be clearly indicated.
	No
	This test is performed at UKHSA Bristol Reference Laboratory.

Hepatitis D (Delta agent)

	Hepatitis D (Delta agent)
	• Antibody detection, • PCR.
	Clotted sample

	10-14 days
	False negatives may occur for a variety of reasons, for example inappropriate timing of sample collection, inappropriate sample, presence of organism below the detectable limit of the assay.
	No
	This test is performed at Virus Reference Department, UKHSA Colindale

Human Herpes Virus-7 DNA Quantitative PCR

For diagnosis of HHV-7 infection.

Human Herpes Virus-7 DNA Quantitative PCR

	False negatives may occur for a variety of reasons, for example inappropriate timing of sample collection, inappropriate sample, presence of organism below the detectable limit of the assay.
	No
	This test is performed at Virus Reference Department, UKHSA Colindale.

Influenza A/B

	Influenza A/B

Joint Fluid Culture

	Joint Fluid Culture
	Gram stain, culture and crystals
	Joint fluid
	Universal (white top)
	Samples should be taken using strict aseptic technique in line with Trust procedure. Ideally a minimum volume of 1ml should be collected. Where adequate sample, also inoculate into blood culture bottle set. Specimens should be sent to the laboratory without delay during normal working hours. Outside of normal working hours samples should be refrigerated. 5

	Collect specimens as per normal procedure. Specimens should be double bagged, with the request form in the outer bag. Specimens should be sent to the laboratory without delay.
	10-14 days
	Suspected cases need to be discussed with the Consultant Microbiologist first.
[redacted]	No.
	This test is performed at RIPL UKHSA Reference Laboratory.

[redacted]e

Mouth Swab Culture

	Negative: 1 –2 days Positive: 2-4 days Culture directed to MRSA only. See Trust MRSA Policy.

	Red topped swab
	No special requirements. Specimens should be sent to the laboratory without delay during normal working hours. Outside of normal working hours samples should be refrigerated.
	10-14 days
	Clinical details are essential. Please state vaccination history.
	No
	This test is performed at Virus Reference Department, Colindale Reference Laboratory

Mycobacterium tuberculosis Fast Track

	M. tuberculosis Fast Track test Detection of *M. tuberculosis*

	Cobas transport media (for urine) Universal (White top)
	No special requirements. Specimens should be sent to the laboratory without delay during normal working hours. Outside of normal working hours samples should be refrigerated.
	10-14 days
	No
	This test is performed at UKHSA Colindale Bacteriology (BRD Colindale) Reference Laboratory.

Neonatal Screen Culture

	Neonatal Screen Culture
	Culture
	• Swabs • Gastric aspirate
	Transport swab (black top) Universal container
	No special requirements. Specimens should be sent to the laboratory without delay during normal working hours. Outside of normal working hours samples should be refrigerated.
	4 days
	Delays in transportation may affect the recovery of pathogens.
	No

Non-indigenous mycoses (e.g., Histoplasma) serology

	Non-indigenous mycoses serology
	Antibody detection

	Specimens should be sent to the laboratory without delay during normal working hours. Outside of normal working hours samples should be refrigerated.
	10-14 days
	Travel history is essential.
	No
	This test is performed at Mycology Reference Laboratory.

Nose Swab Culture

	Nose Swab Culture
	Culture
	Nose swab
	Transport swab (black top) Optimally collected before antimicrobial therapy started. No special requirements.

	• Perianal swab.
	<u>Laboratory no longer supplies collection kits.</u> • Sellotape slide, • Cotton-wool swab in dry container.

Apply clear Sellotape to the perianal region, pressing the adhesive side of the tape firmly against the left and right perianal folds several times; the tape can be wrapped around a tongue depressor to aid specimen collection. Smooth the tape back on the slide, adhesive side down.

Perianal specimens are best obtained in the morning before bathing or defecation. Three specimens should be taken on consecutive days before pinworm infection is ruled out. Cotton-wool swab in dry container should be used for collection.
Spread buttocks apart and rub the moistened cotton wool swab over the area around the anus, but do not insert into the anus. Place cotton wool swab back in its container (no transport medium required).
Occasionally, an adult worm may be collected from a patient and sent in saline or water for identification.
Specimens should be sent to the laboratory without delay during normal working hours.

	Clotted blood
	Yellow top (minimum volume 3.5 ml)
	No special requirements. Specimens should be sent to the laboratory without delay during normal working hours. Outside of normal working hours samples should be refrigerated.
	10-14 days
	Clinical details are essential for processing.
	No
	This test is performed at UKHSA Bristol Reference Laboratory.

Parvovirus B19 PCR

	Parvovirus B19 PCR
	PCR

	Trap bottle Green topped swab Red topped swab
	No special requirements. Specimens should be sent to the laboratory without delay during normal working hours. Outside of normal working hours samples should be refrigerated.
	1 day
	Clinical details are essential for processing
	No

Rubella Serology

	Rubella Serology IgG/IgM
	Antibody detection
	Clotted blood
	Yellow top (minimum volume 3.5 ml)
	No special requirements. Specimens should be sent to the laboratory without delay during normal

	Genotyping, Toxin detection, etc.
	Pure culture
	N/A
	N/A
	10-14 days
	Consultant Microbiologist will decide which isolates need to be sent for further investigation.
	No
	These tests are performed at UKHSA Colindale Bacteriology Reference Laboratory.

Syphilis Serology

	Syphilis antibody detection.
	Antibody detection
	Clotted blood
	Yellow top (minimum volume 3.5 ml)
	No special requirements. Specimens should be sent to the laboratory without delay during normal working hours. Outside of normal working hours samples should be refrigerated.
	4 days, longer if the sample is positive and needs sending to the reference laboratory for confirmation.
	Clinical details are essential for processing.
	No
	Positive in-house samples are sent to UKHSA Bristol Reference Laboratory for full confirmation.

Syphilis IgM, RPR, TPPA, alternative EIA and Immunoblots

	Syphilis IgM, RPR, TPPA, alternative EIA and Immunoblots
	Antibody detection
	Clotted blood
	Yellow top (minimum volume 3.5 ml)
	No special requirements. Specimens should be sent to the laboratory without delay during normal working hours. Outside of normal working hours samples should be refrigerated.
	5-7 days (Longer if sample needs to be sent to Reference Laboratory).
	No
	If sample is positive, it is sent to UKHSA Bristol Reference Laboratory for confirmation.

Syphilis PCR

	Syphilis PCR
	Detection of *Treponema pallidum* DNA

	Swab in viral transport media
	Green topped swab Red topped swab
	No special requirements. Specimens should be sent to the laboratory without delay during normal working hours. Outside of normal working hours samples should be refrigerated.
	10-14 days
	No
	This test is performed at PHE Colindale (STBRU) Reference Laboratory.

TB Culture (Urine)

	TB Culture (Urine)
	Culture
	First-pass early morning urine (from 3 consecutive days)
	Universal (60ml wide-mouth, metal top) 3 specimens should be sent from 3 consecutive days.

Throat Swab Culture

	No
	Confirmation of positive results sent to Swansea Hospital.

Tropical Disease Serology (Regional Travel Screen)

Various including Dengue, Viral haemorrhagic fevers, etc.

	Tropical disease (e.g., Dengue, VHF, etc.) antibody detection
	Antibody detection
	Clotted blood
	Yellow top (minimum volume 3.5 ml)
	No special requirements. Specimens should be sent to the laboratory without delay during normal working hours. Outside of normal working hours samples should be refrigerated.
	7-14 days
	Clinical details including countries visited & dates are essential. Vaccinations & antibiotics given are essential as may affect test results. Contact duty Consultant Microbiologist if required
	No
	This test is performed at PHE Porton Down Reference Laboratory.

Ulcer Swab

	Ulcer Swab
	Culture
	Ulcer swab

	Green top tube with boric acid.
	No special requirements. Specimens should be sent to the laboratory without delay during normal working hours. Outside of normal working hours samples should be refrigerated.
	1 day
	Please contact Laboratory if urgent processing is required.
	No

Urine Culture

	Urine Culture
	Microscopy & Culture
	Urine
	Green top tube with boric acid. Container without boric acid Universal (white top)
	No special requirements. Specimens should be sent to the laboratory without delay during normal working hours.

	urinary tract infection, complicated UTI or those who have not responded to empirical therapy (usually trimethoprim or nitrofurantoin). Delays in transportation may affect the recovery of pathogens. For more detail guidance, please refer to: https://www.gov.uk/government/publications/urinary-tract-infection-uti-diagnosis

Urinary Parasitology (Schistosomiasis)

	Microscopy for *Schistosoma haematobium*
	Microscopy

The sample needs to be collected in the morning. Patient must abstain from sexual activity for 48 hours (but no longer than 7 days) before collecting the sample. Sample should be collected into the provided container (it can be collected from GP or fertility specialist). Sample should be collected by masturbation and the entire specimen should be collected into the container. Please do not use a sheath or condom for collection as they are harmful to sperm. After the collection, please keep the sample close to the body whilst travelling as sperm is sensitive to temperature changes.
To ensure the test results will be accurate, please aim to deliver the sample within 1 hour of its production.

Document Title: Q-Pulse Index:

Revision Number: **Author:** [redacted]

Investigation	Post vasectomy semen analysis
icroscopy	Test

sexual activity for 48 hours (but no longer than 7 days) before collecting the sample. Sample should be collected into the provided container (it can be

The assay is suitable for detection of viruses from nasopharyngeal swabs, nasal swabs, or nasal

[redacted]

For further guidance on handling medicolegal samples and preserving the chain of evidence please refer to this website: New guidance for handling medicolegal samples and preserving the chain of evidence (rcpath.org)

16.4. The Human Tissue Act

Salisbury District Hospital NHS Foundation Trust are licensed by the Human Tissue Act (HTA) to undertake examinations of post-mortem samples submitted by clinical consultants and pathologists. Under the license, the samples may be retained until the examination has been completed and in line with the sample retention policies. It is the obligation of the requesting clinician or pathologist to ensure that examination of samples they submit have been requested by the coroner or appropriate consent has been obtained from the deceased person or their relatives. Only the specific examinations requested by the sending clinician or pathologist may be performed.
If additional work on samples from the deceased is thought necessary by the medical microbiologist or virologist, they must obtain written confirmation of consent from the sending departments. All relevant material is stored securely and under conditions which maintain the integrity of the sample if possible and confidentiality is maintained in compliance with Caldicott principles, as are all samples received. Following processing, relevant material is only retained for the period of time specified by the retention policy.
For further information please refer to Human Tissue Act 2004:
Human Tissue Act 2004 (legislation.gov.uk)

17. Feedback on our microbiology service and complaints procedure

The laboratory is committed to providing a high-quality service to all service users, however, it understands that aspects of the service may not always meet the requirements of the customer. Should this occur and there be a requirement to give feedback or make a complaint to the laboratory please submit your feedback or complaint in writing to the following:

[redacted]	[redacted]
[redacted]	[redacted]
[redacted]	[redacted]